


66TH CONFERENCE ON EXCEPTIONAL CHILDREN

Are Students with Hearing Loss at Risk for Listening-Related Fatigue?

Hilary Davis, Au.D.
Pediatric Audiologist
Vanderbilt Bill Wilkerson Center



SELF-ASSESSMENT: A Journey of Change

PUBLIC SCHOOLS OF NORTH CAROLINA
State Board of Education | Department of Public Instruction

November 8-10, 2016

Disclaimer

Presentation materials are for registered participants of the 66th Conference on Exceptional Children. The information in this presentation is intended to provide general information and the content and information presented may not reflect the opinions and/or beliefs of the NC Department of Public Instruction, Exceptional Children Division. Copyright permissions do not extend beyond the scope of this conference.

SELF-ASSESSMENT: A Journey of Change

66TH CONFERENCE ON EXCEPTIONAL CHILDREN

Acknowledgements

Faculty Investigators

- o Fred H. Bess
- o Ben W.Y. Hornsby
- o Stephen Camarata
- o Alexandra Key

Lab Staff (current and past)

- | | | |
|----------------------|-----------------------|------------------|
| o Nicholas Bennett | o Emily Fustos | o Quela Royster |
| o Angela Chen | o Elizabeth Geller | o Rebecca Schoon |
| o Tonia Davis | o Amanda Headley | o Sara Seckman |
| o Stone Dawson | o Dorita Jones | o Amelia Shuster |
| o Andy DeLong | o Ralph Leverett | o Beth Suba |
| o Caralie Focht | o Ronan McGarrigle | o Maureen Virts |
| o Samantha Gustafson | o Lindsey Rentmeester | o Ye Wang |
| o Olivia Gutierrez | o Virginia Rich | o Krystal Wertel |

SELF-ASSESSMENT: A Journey of Change

66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

Today's Goals

- Definitions of listening effort and fatigue
- Consequences of fatigue
- How do you measure fatigue?
 - Behavioral, physiologic, subjective


SELF-ASSESSMENT:   

A Journey of Change




66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

What is Fatigue?



- Affects several areas of life including *physical, emotional, and cognitive or mental* domains
- **Physical fatigue:** reduced ability or desire to perform some physical task
- **Cognitive/mental fatigue:** feeling of tiredness, exhaustion, or lack of energy due to cognitive or emotional demands

SELF-ASSESSMENT:   

A Journey of Change

66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER




Aren't we all fatigued?

Fatigue is one of the most common complaints reported in primary care settings

- Transient fatigue is common, even in healthy populations

Recurrent, severe fatigue

- Uncommon in healthy populations, but common in many chronic health conditions
 - Previous reports in individuals with cancer, HIV AIDs, Parkinson's, Multiple Sclerosis


SELF-ASSESSMENT:   





A Journey of Change

66TH CONFERENCE ON EXCEPTIONAL CHILDREN
Slide from Gustafson, 2015

VANDERBILT UNIVERSITY
MEDICAL CENTER


“When you are hard of hearing you struggle to hear; when you struggle to hear you get tired; when you get tired you get frustrated; when you get frustrated you get bored; when you get bored you quit.” (Pichora-Fuller, 2003)







SELF-ASSESSMENT:    
A Journey of Change 66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

Listening in the Classroom



SELF-ASSESSMENT:    
A Journey of Change 66TH CONFERENCE ON EXCEPTIONAL CHILDREN





VANDERBILT UNIVERSITY
MEDICAL CENTER

Listening in the Classroom

- Degraded signal from effect of hearing loss and poor classroom acoustics
- Does the child put forth additional EFFORT to listen and understand in their typical listening setting? (i.e. classroom, cafeteria, gymnasium, after-school activities, etc)

CHL and AHL must increase mental effort compared to those without HL when attempting to detect, process, and respond to auditory stimuli (Hicks and Tharpe, 2002; McCoy et al., 2005)

- Increase in LISTENING EFFORT (Hornsby, 2013)

SELF-ASSESSMENT:    
A Journey of Change 66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

Does effortful listening affect fatigue?

Figure 1. Conceptual Model Linking Hearing Loss to Fatigue and School Performance. Shaded areas represent events that occur repeatedly throughout the school day.

Bess & Hornsby, 2013

SELF-ASSESSMENT: **A Journey of Change** 66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

What does a CHL feel at the end of a school day? Is their ability to perform at school affected?

SELF-ASSESSMENT: **A Journey of Change** 66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

Why do we care? Negative Consequences of Fatigue

ADULTS	CHILDREN
Report stress, inattention, concentration difficulties	Higher rates of absenteeism at school
Reduced mental processing and decision-making capabilities	Poorer school achievement
Less productive and more prone to accidents at work	Difficulties with attention, concentration, and distractibility
Less active and more isolated	More likely to fail a grade

SELF-ASSESSMENT: **A Journey of Change** 66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

Measuring Listening-Related Fatigue

Subjective Measures

- Questionnaires, rating scales that query an individual's mood or feelings
 - Discussed in Part 2

Objective Measures

- Does not require the individual to make a judgment about their fatigue
 - Behavioral: measuring decrement in performance
 - Physiological: measuring physiologic changes associated with mental effort
 - Cortisol
 - EEG

SELF-ASSESSMENT:
A Journey of Change

66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

Vanderbilt Study on Listening Effort & Fatigue

- 6-12 year old children
 - Bilateral, mild to moderately-severe, permanent hearing loss
- Inclusion/Exclusion:
 - No cochlear implant users
 - General education classroom
 - Monolingual English speakers
 - No diagnosis of cognitive impairment, autism, or other developmental disorder
- Experimental group (n=59)
 - 30 males, 29 females
 - Age = 9.93 (1.92) years
- Control Group (n=42)
 - 25 males, 17 females
 - Age = 9.16 (2.32) years

The graph plots Hearing Level (dB HL) on the y-axis (from -20 to 120) against Frequency (Hz) on the x-axis (logarithmic scale from 125 to 8000). Three data series are shown: Right Ear (open circles), Left Ear (filled circles), and Audiometer Limits (asterisks). The hearing levels for both ears are generally between 20 and 40 dB HL across the frequency range, with some variability at higher frequencies. The audiometer limits are shown as a dashed line at approximately 120 dB HL.

Gustafson, 2015

SELF-ASSESSMENT:
A Journey of Change

66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

Visit Descriptions

- Initial Visit
 - Qualifying testing including non-verbal IQ and language measures
- Experiment 1: Cortisol
- Experiment 2/3: Simulating Fatigue (ERP and dual-listening tasks)*
- Experiment 4: Language Testing

*completed in both aided and unaided conditions in CHL


SELF-ASSESSMENT:
A Journey of Change

66TH CONFERENCE ON EXCEPTIONAL CHILDREN
Slide from Gustafson, 2015


VANDERBILT UNIVERSITY
MEDICAL CENTER

Research Question #1

- Do children with mild to moderate sensorineural hearing loss exhibit increased fatigue (as measured by cortisol responses) throughout the course of a typical school day compared to their normal hearing counterparts? If so, how does the cortisol response differ?



CORTISOL

SELF-ASSESSMENT:  **A Journey of Change** 66TH CONFERENCE ON EXCEPTIONAL CHILDREN


VANDERBILT UNIVERSITY
MEDICAL CENTER

Cortisol Patterns in Children with Hearing Loss

Hicks and Tharpe (2002)


- Children with mild to moderate hearing loss (n=10) and a control group (n=10)
- 5-11 years old
- Salivary cortisol at 9am and 2pm

No significant difference in cortisol levels between children with hearing loss and controls



Group	Morning	Afternoon
Hearing Loss	~0.40	~0.25
Normal Hearing	~0.35	~0.22


Figure adapted from original publication by Gustafson, 2015

SELF-ASSESSMENT:  **A Journey of Change** 66TH CONFERENCE ON EXCEPTIONAL CHILDREN


VANDERBILT UNIVERSITY
MEDICAL CENTER

Stress, Cortisol, and Fatigue

- Stress is the body's reaction to change that requires a physical, mental, or emotional response
 - Stress is caused by good and bad experiences
- Cortisol** levels provide a physiologic measure of stress
 - Regulated by the hypothalamic-pituitary-adrenal (HPA) axis
 - Related to sugar levels in the blood that fluctuate based on the need to mobilize energy




Gustafson, 2015

SELF-ASSESSMENT:  **A Journey of Change** 66TH CONFERENCE ON EXCEPTIONAL CHILDREN

“Typical” Cortisol Patterns

In non-fatigued individuals, cortisol levels have a typical diurnal pattern

- Build-up of cortisol during sleep
 - Rapid rise upon awakening
 - Cortisol Awakening Response; CAR
 - Slow decline in cortisol throughout the day
- Jerjes, et al., (2005).




Gustafson. 2015

SELF-ASSESSMENT:
A Journey of Change

46TH CONFERENCE ON EXCEPTIONAL CHILDREN

“Atypical” Cortisol Patterns

- Sustained stress or fatigue can lead to abnormal diurnal cortisol patterns
 - Reduced response with “Chronic Fatigue Syndrome”
-
- Salivary cortisol (nmol/L)
- Time (h)
- Controls
- CFS
- Legend: ● Controls, ○ CFS
- | Time (h) | Controls (nmol/L) | CFS (nmol/L) |
|----------|-------------------|--------------|
| 08:00 | ~14 | ~8 |
| 12:00 | ~10 | ~6 |
| 16:00 | ~6 | ~5 |
| 20:00 | ~2 | ~4 |



Jerjes, et al., (2005).


Gustafson, 2015

SELF-ASSESSMENT:
A Journey of Change

66TH CONFERENCE ON EXCEPTIONAL CHILDREN

“Atypical” Cortisol Patterns

- Sustained stress or fatigue can lead to abnormal diurnal cortisol patterns
 - Reduced response with “Chronic Fatigue Syndrome”
 - “Elevated” CAR in patients with depression
-
- | Time | Healthy (nmol/l) | Depression (nmol/l) | CFS (nmol/l) |
|-------|------------------|---------------------|--------------|
| 08:00 | ~10 | ~10 | ~10 |
| 12:00 | ~25 | ~20 | ~20 |
| 16:00 | ~40 | ~35 | ~25 |
| 20:00 | ~35 | ~25 | ~20 |



Bhagwagar, Z., Hafizi, S., & Cowen, P. J. (2005).

Gustafson, 2015

SELF-ASSESSMENT:
A Journey of Change

66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER




Vanderbilt Study: Cortisol Measurement

- Participants
 - Children with hearing Loss (n=32)
 - Control group (n=28)
- Six samples per day


1. Awakening*	4. 10:00 am
2. 30 min post-wake up*	5. 2:00 pm
3. 60 min post-wake up*	6. 8:00 pm*
- Sampled on two separate school days


*Samples taken by parents at home


Bess, Gustafson, Corbett, Lambert, Camarata, and Hornsby (2016)


SELF-ASSESSMENT:   
A Journey of Change


66TH CONFERENCE ON EXCEPTIONAL CHILDREN












VANDERBILT UNIVERSITY
MEDICAL CENTER

Vanderbilt Study on Listening Effort & Fatigue

Cortisol Findings:
Diurnal cortisol patterns in children with hearing loss are not "typical"

- Elevated levels at awakening and reduced CAR may suggest increased stress
 - Similar to adults with high "burnout"
 - Indicative of a dysregulation in HPA-axis activity


Gustafson, 2015

SELF-ASSESSMENT:  *A Journey of Change* 66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

Research Question #2

Will hearing-related fatigue for CHL differ from the normal hearing children as measured by ERPs?

SELF-ASSESSMENT:  *A Journey of Change* 66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

Measuring Fatigue with ERP


Event-related potentials (ERP) are changes in ongoing EEG activity that are time-locked to the onset of the auditory event

- Reflects change in brain activity associated with the processing of that stimulus


Centro-parietal P300 response

- Sensitive to fatigue due to cognitive processing (Murata, Uetake, & Takasawa, 2005; Uetake & Murata, 2000)

More fatigue → reduced amplitude



Gustafson, 2015

SELF-ASSESSMENT:  *A Journey of Change* 66TH CONFERENCE ON EXCEPTIONAL CHILDREN


VANDERBILT UNIVERSITY
MEDICAL CENTER

Assessing Fatigue in the Lab




































































































3 hours


ERP 1 → Speech Processing Tasks → ERP 2


- Participants
 - Children with hearing Loss (n=33)
 - Control group (n=29)
- Stimuli
 - Oddball paradigm (70/30)
 - Speech syllables ("gi" and "gu") at 65 dB SPL
 - Multi-talker babble at +10 dB SNR
- Procedure
 - 128-channel Geodesic sensor net
 - Passive task




Gustafson, 2015

SELF-ASSESSMENT:                                                                                                   







VANDERBILT UNIVERSITY
MEDICAL CENTER

Who's at risk?

Children with Hearing Loss

$r = .114$

Reduced Cognitive Processing

This lack of relationship with degree of hearing loss is consistent with subjective fatigue data and suggests that children with even mild hearing loss are at increased risk for fatigue.

Age, Language, Nonverbal Intelligence, and Speech in Noise Recognition did not significantly account for variability in cognitive processing changes associated with listening-related fatigue.

Gustafson, 2015

SELF-ASSESSMENT:

A Journey of Change 66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

Vanderbilt Study on Listening Effort & Fatigue

Findings:

- Laboratory testing can induce listening-related fatigue.
 - Auditory-evoked P300 can be used to measure changes in cognitive processing associated with listening-related fatigue.
- Compared to children with normal hearing, children with mild-to- moderately-severe hearing loss:
 - demonstrate delayed cognitive processing time during active discrimination of speech in babble noise.
 - show similar consequences (i.e., reduced cognitive processing) of speech-processing related fatigue.

Gustafson, 2015

SELF-ASSESSMENT:

A Journey of Change 66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

Answers to Research Questions

- Do children with mild to moderate sensorineural hearing loss exhibit increased fatigue (as measured by cortisol responses) throughout the course of a typical school day compared to their normal hearing counterparts? If so, how does the cortisol response differ?**
 Yes, CHL have "atypical" cortisol patterns, suggesting increased stress.
- Will hearing-related fatigue for CHL differ from the normal hearing children as measured by ERPs?**
 Yes, CHL present show delays in cognitive processing of speech in noise discrimination. Both children with and without HL show reductions in cognitive processing secondary to speech-processing related fatigue.

SELF-ASSESSMENT:


A Journey of Change 66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

CHL at Risk for Fatigue?

YES!


So what do we do about it?

SELF-ASSESSMENT:  *A Journey of Change* 66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

Stick around for PART 2!

- Discussing subjective listening-related fatigue, commonly reported symptoms at home and in the classroom, and potential management strategies for CHL.

SELF-ASSESSMENT:  *A Journey of Change* 66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

Questions? Thoughts?



SELF-ASSESSMENT:  *A Journey of Change* 66TH CONFERENCE ON EXCEPTIONAL CHILDREN

VANDERBILT UNIVERSITY
MEDICAL CENTER

Visit the Listening and Learning Lab's website at
<http://my.vanderbilt.edu/listeninglearninglab>
Hilary.Davis@Vanderbilt.edu



SELF-ASSESSMENT:  
A Journey of Change

66TH CONFERENCE ON EXCEPTIONAL CHILDREN
